

SEPSIS-ADULT

I. **PURPOSE:** To facilitate rapid identification and treatment of patients with suspected sepsis.

II. FIELD ASSESSMENT/TREATMENT INDICATORS

For possible infection and two or more of the following:

- Temperature > 100.4 F or < 96.8 F
- Sustained HR >90
- EtCO2 < 25 mmHg
- Sustained Respiratory rate >20

III. BLS INTERVENTIONS

- Oxygen therapy as clinically indicated.
- Position patient as tolerated and cover to avoid shivering. If altered, place in left lateral position.
- Obtain and assess blood glucose level. If indicated, administer Glucose Oral per ICEMA Reference #11010 Medication Standard Orders.

IV. LIMITED ALS INTERVENTIONS

- Perform activities identified in the BLS Interventions
- Obtain vascular access.
- Administer 500 ml IV bolus, may repeat one (1) time.
- If hypoglycemic;
 - > Dextrose per ICEMA Reference #11010 Medication Standard Orders, or
 - If unable to establish IV, Glucagon may be given one (1) time per ICEMA Reference #11010 - Medication - Standard Orders.
- If hyperglycemic;
 - Administer 500 ml IV bolus may repeat one (1) time.

V. ALS INTERVENTIONS

- Perform activities identified in the BLS and LALS Interventions.
- Place on cardiac monitor and obtain a 12-lead ECG as clinically indicated.
- Monitor EtCO2

For profound hypotension, unresponsive to fluid boluses, administer Push Dose Epinephrine per ICEMA Reference #11010 - Medication - Standard Orders

VI. SPECIAL CONSIDERATIONS

- Risk Factors:
 - Indwelling catheters (foley, PICC line etc.)
 - Recent surgery
 - > Open wounds
 - Bedridden or immobile patients
 - Compromised Immune system due to comorbidities (cancer, autoimmune disease, etc.)
- Prioritize early recognition, IV fluids and rapid hospital notification and transport.
- If signs of pulmonary edema, stop or limit fluid boluses.
- Hypotension is a late indicator for septic shock.

VII. REFERENCES

<u>Number</u>	<u>Name</u>
11010	Medication - Standard Orders
11020	Procedure - Standard Orders